

# Fine mapping *SPP1*, a QTL controlling the number of spikelets per panicle, to a BAC clone in rice (*Oryza sativa*)

Touming Liu · Donghai Mao · Shengpeng Zhang ·  
Caiguo Xu · Yongzhong Xing

Received: 26 June 2008 / Accepted: 17 February 2009 / Published online: 6 March 2009  
© Springer-Verlag 2009

**Abstract** Near isogenic lines (NILs) can be used to efficiently handle a target quantitative trait locus (QTL) by blocking genetic background noise. One QTL, *SPP1*, which controls the number of spikelets per panicle (SPP), was located on chromosome 1, near *Gn1a*, a cloned gene for rice production in a recombinant inbred line population. NILs of the *SPP1* regions were quickly obtained by self-crossing recombinant inbred line 30 which is heterozygous around *SPP1*. Using a random NIL-F<sub>2</sub> population of 210 individuals, we mapped *SPP1* to a 2.2-cM interval between RM1195 and RM490, which explained 51.1% of SPP variation. The difference in SPP between the two homozygotes was 44. F<sub>2</sub>-1456, one NIL-F<sub>2</sub> plant, was heterozygous in the *SPP1* region but was fixed in the region of *Gn1a* gene. This plant F<sub>3</sub> family showed a very wide variation in SPP, which suggested that it was *SPP1* but *Gn1a* affected the variation of SPP in this population. In a word, *SPP1* is a novel gene distinct from *Gn1a*. Four newly developed InDel markers were used for high-resolution mapping of *SPP1* with a large NIL-F<sub>2</sub> population. Finally, it was narrowed down to a bacterial artificial chromosome clone spanning 107 kb; 17 open reading frames have been identified in the region. Of them, LOC\_Os01g12160, which encodes an IAA synthetase, is the most interesting candidate gene.

## Introduction

Grain yield is one of the most valuable traits in crop production. Pursuing high grain yield is one of the most important goals in rice (*Oryza sativa* L.) production. Grain yield is frequently replaced with grain yield per plant in quantitative trait locus (QTL) mapping owing to control of field experimental scale (Zhuang et al. 2002; Hittalmani et al. 2003). Grain yield per plant is determined by its three components, panicles per plant, 1,000-grain weight, and spikelets per panicle (SPP). Among the three components, SPP frequently makes the greatest contribution to grain yield and has received the most attention in genetic analysis. However, this trait is inherited in a quantitative manner and typically controlled by a number of major and minor QTL and is also affected by environment, which present a challenge in characterizing SPP. With QTL analysis based on molecular markers, SPP has been studied, and hundreds of QTL have been reported (Yu et al. 1997; Zhuang et al. 1997; Xing et al. 2002; Septiningsih et al. 2003). Many reports have provided evidence for a very complicated genetic basis of yield traits, which are affected simultaneously by QTL, epistasis, and environment in primary mapping populations (Li et al. 1997; Xing et al. 2002; Zhuang et al. 2002). Currently, QTL fine mapping and cloning are still difficult although many QTL for SPP have been mapped to chromosome regions.

In a primary mapping population with simultaneous segregation of multiple genetic factors, a given trait value is contributed not only by a given QTL but also by other QTL. For fine mapping of a QTL, genetic background noise should be avoided. Hence, QTL analysis in advanced populations is necessary for a good understanding of their characteristics. Near isogenic lines (NILs) has been proven to be an ideal population for QTL fine mapping and cloning.

---

Communicated by H. H. Geiger.

---

T. Liu · D. Mao · S. Zhang · C. Xu · Y. Xing (✉)  
National Key Laboratory of Crop Genetic Improvement  
and National Center of Plant Gene Research (Wuhan),  
Huazhong Agricultural University, 430070 Wuhan, China  
e-mail: yzxing@mail.hzau.edu.cn; yzhxing@hotmail.com

Many QTL have been fine mapped and cloned using a large NIL population in which a major QTL can be visualized as a Mendelian factor (Ren et al. 2005; Song et al. 2007; Xing et al. 2008; Xue et al. 2008). There are three strategies for QTL-NILs construction. Consecutive backcrossing (CB-NILs) combined with marker aided selection (MAS) is the conventional method for developing NILs and has been widely used (Yamamoto et al. 1998, 2000; Li et al. 2004; Fan et al. 2006). However, CB-NILs construction is time consuming; usually taking 3 years to perform in rice after QTL information is obtained. The alternative method is to search directly for NILs according to trait performance (TP-NILs) in advanced recombinant inbred lines (RILs) (Zhang et al. 2006). Its genetic principle is that only a very small portion of the genomic regions of a RIL (e.g.,  $F_6$  or  $F_7$ ) segregates, and if the segregating region contains a major QTL, trait variation would be exhibited within the RIL. Compared with CB-NILs development, TP-NILs development is faster. In most cases, however, TP-NILs is effective for only major QTL because variation of the trait controlled by major QTL is easily observed. The third strategy for NILs' development is the heterogeneous inbred family (HIF-NILs) method (Tuinstra et al. 1997), which is based on molecular marker screening for a small heterozygous genomic region. HIF-NILs' development is effective for the target QTL whether it is a major or minor QTL. The genetic principle of TP-NILs and HIF-NILs is similar. Indeed, both TP-NILs and HIF-NILs result in the selection of a RIL that is heterozygous in a small genomic region, but TP-NILs selection is based on a phenotypic screening, whereas HIF-NILs selection is based on a molecular marker screening.

Recently, more and more rice SPP QTL are being fine mapped and cloned using NILs. In NIL- $F_2$ , two QTL, *Gn1a* and *Gn1b*, were detected in the QTL *Gn1* region, and *Gn1a* was fine mapped to a 6.3-kb region. Gene cloning analysis shows that *Gn1a* is a gene for cytokinin oxidase/dehydrogenase (OsCKX2) which controls the grain number phenotype variation (Ashikari et al. 2005). *gpa7*, a QTL controlling SPP on chromosome 7, was narrowed down to a 35-kb region that contains five predicted genes (Tian et al. 2006). *qSPP7*, a major QTL for SPP, was fine mapped to a 0.2-cM region (Xing et al. 2008), renamed *Ghd7*, and finally cloned (Xue et al. 2008).

With the complete genome sequence of rice, chromosome-walking will be greatly reduced or even avoided in the process of map-based gene isolation, and the Rice Genome Automated Annotation System (<http://RiceGAAS.dna.affrc.go.jp>), combined with the high-resolution of gene fine mapping, is helpful in identifying the candidate gene. Now that thousands of QTL have been mapped during the last two decades, the focus of QTL research is being gradually transferred to QTL isolation from mapping in rice.

In our previous study, a population consisting of 190 RILs was developed for QTL mapping by single-seed descents from a cross between two *indica* cultivars, Zhen-shan 97 (ZS97), a widely used maintainer for hybrid production, and Teqing (TQ), a high yielding variety in China. One major QTL, *SPP1*, was detected in the interval between RM490 and RM283 near *Gn1a* and *Gn1b* on chromosome 1. *SPP1* had an additive effect of 16.5 and 15.4 in two environments, respectively. Because of low resolution of QTL mapping in RILs, the relationship between *SPP1* and *Gn1a* and *Gn1b* is not clear. To characterize *SPP1*, we used in this study a straightforward HIF strategy to develop a NIL population and fine map it to a bacterial artificial chromosome (BAC) clone with a large NIL population.

## Materials and methods

### Experimental populations and phenotypic measurements

In the mapping population consisting of 190 RILs, the line RIL30 is homozygous at all investigated 176 marker loci except for the two markers RM490 and RM283, which flank the QTL *SPP1*. Naturally, selfing progenies of RIL30 (pseudo  $F_2$ ) constitute a NIL- $F_2$  population of *SPP1*. In the summer of 2005, a total of 2,200 NIL- $F_2$  plants were planted into a bird-net-equipped field in the experimental farm at Huazhong Agricultural University, Wuhan, China. Ten seedlings (approximately 25 days old) per row were transplanted with a distance of 16.5 cm between plants within a row, and 26.4 cm between rows. A random  $F_2$  population of 210 plants (subpopulation) from the 2,200 plants was used for QTL analysis. In May 2006,  $F_3$  families of the 210  $F_2$  plants; the progeny of  $F_2$ -1456, which is a NIL- $F_2$  plant segregated at only *SPP1* but fixed at *Gn1a*; and the progeny of five recombinants (1208, 1367, 1599, 1758 and 1747) were sowed on a seedling bed. For QTL analysis on progeny test, 24 seedlings, approximately 25 days old, of each family were transplanted into the same experimental farm with the same planting density as that of NIL- $F_2$ . Because of bad germination of 44 families, 166 families were used. 240  $F_3$  seedlings of  $F_2$ -1456 were transplanted into a 20-row plot. Sixty  $F_3$  seedlings of each recombinant were transplanted into a five-row plot. Progenies from two homozygous  $F_2$  with ZS97 alleles and TQ alleles at *SPP1* were grown as the controls (CK-Z and CK-T, respectively). At the ripening stage, all 2,200 NIL- $F_2$  plants, 20 plants in the middle of the two rows of each NIL- $F_3$ , 180 plants except the plants seated in the boundaries of  $F_3$ -1456, and 50 plants except the plants seated in the boundaries of each recombinant were harvested individually to score the following traits: SPP as the total number of spikelets from the whole plant divided by its number

of tillers, the yield per plant as the total weight (g) of the grains from the entire plant, and the number of filled grains per panicle as the total number of grains from the entire plant divided by the number of tillers. The heading date of each plant was scored as the days from sowing to the first appearance of a panicle.

#### DNA extraction and molecular marker development

DNA was extracted from fresh leaves at the seedling stage using the CTAB method (Murray and Thompson 1980). Simple sequence repeat (SSR) markers in the *SPP1* region were identified from the Gramene database (<http://www.gramene.org/>). The SSR primers were designed according to the public database (Temnykh et al. 2000, 2001; McCouch et al. 2002). Four newly developed InDel markers are listed in Table 1 on the basis of the available public rice genome sequences ([http://www.rgp.dna.affrc.go.jp](http://www.rgp.dna.affrc.go.jp;); <http://www.rise.genomics.org.cn/>).

#### Data analysis

SSR marker assay was conducted as described by Wu and Tanksley (1993). The local molecular linkage map was constructed on the basis of the subpopulation of 210 plants using Mapmaker 3.0 (Lincoln et al. 1992) and the Kosambi function to calculate the genetic distance. Interval QTL mapping was performed with both  $F_2$  and  $F_3$  data with the use of Mapmaker/QTL 1.1 (Lander and Botstein 1989; Lincoln et al. 1993). In the progeny test, the arithmetic mean of each family was regarded as its phenotype value for QTL analysis.

#### Putative ORF prediction in 107 kb region

The putative open reading frame (ORF) in the 107-kb region was predicted by the online software FGESH on the basis of the gene structure available on the Internet (<http://linux1.softberry.com/berry.phtml?topic=fgesh&>

[group=programs&subgroup=gfind](#)), and monocot plants were set as the model organism for prediction.

## Results

#### Trait variation in the subpopulation

The two parents, Zhenshan 97 (ZS97) and Teqing (TQ), showed highly significant differences in SPP. ZS97 exhibited a small panicle of 91 spikelets, and TQ had a large panicle of more than 226 spikelets. The SPP showed a normal distribution whose variation range in  $F_2$  was 129.4–243.5 (Fig. 1). However, a progeny test confirmed that 45 and 41 families expressed identical large and small SPP, respectively, whereas 80 families showed varied SPP. Frequencies of the three genotypes was fitted to the expected Mendelian ratio (1:2:1) for single locus segregation ( $\chi^2 = 0.4$ ,  $P = 0.81$ ). This analysis suggested that one QTL/gene controlled SPP in the  $F_2$  population. In the  $F_2$  subpopulation, no plant had an SPP less than the mean of ZS97 SPP minus twofold its standard deviation, and more than the mean of TQ SPP plus twofold its standard deviation. This finding indicates that no transgressive segregation was observed for SPP in either direction and that the QTL did not express over dominance. Meanwhile, the subpopulation headed within 1 week.

#### QTL analysis for *SPP1*

Five markers inside the region from RM283 to RM490 were used to genotype the  $F_2$  population from RIL30 self-crossing, and a local linkage group covering 11.6 cM was constructed (Fig. 2). QTL analysis based on the  $F_2$  population defined the *SPP1* to a 2.2-cM region between the markers RM1195 and RM490, which was closely linked to *Gn1a* flanked by RM1195 and RM3604 markers (Ashikari et al. 2005) (Fig. 2). The QTL explained 51.1% of phenotype variance with additive effect of 22 spikelets (Table 2).

**Table 1** InDel markers developed for fine mapping of the *SPP1* locus


| Marker | InDel size (bp) <sup>a</sup> | Size (bp) <sup>b</sup> | Forward primer (5'–3') | Reverse primer (5'–3') | Physical position                           |
|--------|------------------------------|------------------------|------------------------|------------------------|---|
| YN23   | 23                           | 258                    | CCGATATCACCTGCTTGGAG   | AGACTAACTGGACGGCGATG   | AP001633 <sup>c</sup> , 121880 <sup>d</sup> |
| YN27   | 6                            | 192                    | TACCACTGAACCCACGTGTC   | GCTGCCTTTGTTCTCACGTT   | AP002094, 3900                              |
| YN34   | 12                           | 348                    | TGTGCCATATCTGTGCCTGT   | TCGGTTAGGTCGGTTAATTCC  | AP002094, 111761                            |
| YN37   | 10                           | 198                    | GACCTCCCTCCTATCGTAATGA | CAAGGAGGGAAGGACAACAA   | AP002094, 57860                             |

<sup>a</sup> InDel size difference between Nipponbare and 93-11 sequence

<sup>b</sup> The size of amplicon in Nipponbare

<sup>c</sup> The accession number of BAC clone where marker located

<sup>d</sup> The marker position in bp in corresponding BAC clone



**Fig. 1** Frequency distribution of number of spikelets per panicle in NIL- $F_2$  population. The three genotypes of homozygous Teqing (TT) and Zhenshan 97 (ZZ) and heterozygote (TZ) at *SPP1* were identified by progeny test


Meanwhile, QTL explaining 24.5 and 13.6% of phenotypic variance of grains per panicle (i.e., +21.4 grains per panicle) and total grain yield (i.e., +5.7 g per plant), respectively, were detected in the same interval, which suggest that the same gene is involved both in SPP and grains per

panicle. However, no QTL affecting tillers per plant and 1,000-grain weight, two components of yield, was detected in this region. In the  $F_3$  progeny test, the QTL could explain 63.4% of SPP variation, which explains more of the trait variation than in the  $F_2$  population (Table 2). Similarly, the LOD value of the QTL for SPP detected in the  $F_3$  progeny is higher than that in the  $F_2$  population, but the additive effect is less than that of the  $F_2$  population. *SPP1* acted as additive in both populations (Table 2). The position of the LOD peak for *SPP1* was 0.2 cM from RM490. No QTL was detected for heading date.

### *SPP1* distinct from *Gn1a* and *Gn1b*

*SPP1* was genetically located in the region near *Gn1a* and *Gn1b* in the subpopulation. To clearly confirm that *SPP1* in this study was not the same as *Gn1a* or *Gn1b*, we investigated SPP variation using the progeny of  $F_2$ -1456, which segregates in the interval between RM8111 and RM490 but fixed in the other intervals containing *Gn1a* and *Gn1b*. A very wide variation of 124.3–233.5 in SPP was very similar

**Fig. 2** The local genetic linkage map and high-resolution physical map for *SPP1* region numbers in brackets on the right/left side of markers indicate the number of recombinants between *SPP1* and the respective markers. The short horizontal lines represent BAC/PAC clones of cv. Nipponbare with their accession numbers (<http://www.gramene.org/>). The dashed lines indicate the relative position of the corresponding marker on BAC/PAC clones



**Table 2** *SPP1* effects on SPP, grains per panicle, and yield per plant in the subpopulation and its progenies

| Trait                 | Generation  | QTL position <sup>a</sup> | LOD   | A <sup>b</sup> | D                    | Var (%) <sup>d</sup> |
|-----------------------|-------------|---------------------------|-------|----------------|----------------------|----------------------|
| Spikelets per panicle | $F_2$       | RM1195 + 2.0 cM           | 23.95 | 22.05          | 0.11 Ns <sup>c</sup> | 51.1                 |
|                       | $F_3$       | RM1195 + 1.6 cM           | 35.52 | 10.81          | 0.55 Ns              | 63.4                 |
| Grains per panicle    | $F_2$       | RM1195 + 2.0 cM           | 10.0  | 21.4           | 9.6                  | 24.5                 |
|                       | $F_3$       | RM1195 + 1.6 cM           | 6.65  | 5.27           | 1.09                 | 19.8                 |
| Yield per plant       | $F_2$       | RM1195 + 2.0 cM           | 5.5   | 5.7            | 0.7 Ns               | 13.6                 |
|                       | $F_3$       | RM1195 + 1.6 cM           | 7.03  | 1.2            | 0.35 Ns              | 19.4                 |
| Spikelets per panicle | $F_3$ -1456 | RM1195 + 1.9 cM           | 12.40 | 21.53          | 2.16 Ns              | 37.0                 |

<sup>a</sup> Position of QTL means the LOD peak position of QTL

<sup>b</sup> Additive effect; positive additive effect indicates that the Teqing allele increased the trait value

<sup>c</sup> Ns non significance at  $P = 0.05$

<sup>d</sup> Percentage of total phenotypic variance explained by the QTL

to that in  $F_2$  population. Averaged SPP of TT and ZZ genotypes were 204.8 and 161.4 with standard deviations of 21.3 and 21.0, respectively. One QTL was still identified in the population, which explains 37% of the phenotypic variance (Table 2). This finding suggested that neither *Gn1a* nor *Gn1b* controlled SPP variation in this population. Their physical positions are illustrated in Fig. 2. *Gn1a* was located about 880 kb from RM1195, which was about 1.2 Mb from *SPP1* (Fig. 2).

#### *SPP1* fine mapping

For fine mapping of *SPP1*, 2,200 selfing progenies of RIL30 were grown. Only 250 plants (less than one-eighth) with the smallest panicle of less than 150 spikelets, which were assumed to be Zhenshan 97 homozygotes at *SPP1*, were used for recombinant screening with markers RM1195 and RM490. Eight and two recombinants were identified between RM1195 and *SPP1*, and RM490 and *SPP1*, respectively. To enhance the resolution of the *SPP1* local linkage map, we used one SSR (RM8111) and four InDel markers (Table 1). These five markers were further used to screen the ten recombinants: six recombinants were identified between YN23/RM8111 and *SPP1*, four between YN27 and *SPP1*, and one between YN34 and *SPP1* (Fig. 2). YN37 was found to co-segregate with *SPP1* (Fig. 2). In case the trait measurement yielded false Zhenshan 97 homozygous plants, progeny tests of the five recombinants (1208, 1367, 1599, 1758, and 1747) between YN27 and YN34 were conducted. Each recombinant progeny showed an identical small SPP, which was highly significantly smaller than the control Teqing homozygotes, but showed no difference with the control Zhenshan 97

homozygotes (Table 3). This result confirmed the Zhenshan 97 homozygous identity of the five recombinants at *SPP1*. Therefore, *SPP1* was narrowed down to the region of approximately 107 kb bounded by YN27 and YN34 (Fig. 2). One Nipponbare BAC (AP002094) exactly covered the region.

#### Putative genes in 107-kb region

There are 17 predicted genes in the 107-kb region according to the rice genome automated annotation database (<http://www.tigr.org/tdb/e2k1/osa1/irgsp.shtml>) (Table 4). Of these, 11 genes have homology with rice full-length cDNAs or expressed sequence tags (ESTs). Among these 17 genes, 11 are of unknown function, one gene is a transposon protein gene, and the functional annotations of the remaining five genes are as follows: LOC\_Os01g12020 is a putative nonspecific lipid-transfer protein (LTP) precursor gene; both LOC\_Os01g12030 and LOC\_Os01g12070 are putative endoglucanase 1 precursor genes; and LOC\_Os01g12130 and LOC\_Os01g12160 are putative MtN3 (*Medicago truncatula* nodulate gene 3) protein-like genes and indole-3-acetic acid-amido synthetase GH3.3 genes, respectively (Table 4). Accordingly, 17 genes were predicted with the use of the online software FGENESH in the 107-kb region (<http://www.softberry.com>). Of these, seven genes (Loc\_Os01g12010, Loc\_Os01g12030, Loc\_Os01g12040, Loc\_Os01g12060, Loc\_Os01g12070, Loc\_Os01g12140, and Loc\_Os01g12150) are completely identical, and the other ten genes have a slight difference in either their gene framework or a different encoding sequence between the putative genes, as predicted by two different methods (Table 4).

**Table 3** SPP performance in the progenies of recombinants

| Recom binants     | Spikelets per panicle |                               | Marker |      |      |      |      |
|-------------------|-----------------------|-------------------------------|--------|------|------|------|------|
|                   | $F_2$ plant           | $F_2$ progenies               | RM8111 | YN23 | YN27 | YN37 | YN34 |
| 1208              | 135.60                | 149.1 $\pm$ 15.5 <sup>c</sup> | TZ     | TZ   | TZ   | ZZ   | ZZ   |
| 1367              | 144.13                | 150.9 $\pm$ 12.9 <sup>c</sup> | TZ     | TZ   | TZ   | ZZ   | ZZ   |
| 1599              | 148.25                | 150.9 $\pm$ 21.2 <sup>c</sup> | TZ     | TZ   | TZ   | ZZ   | ZZ   |
| 1758              | 141.00                | 145.7 $\pm$ 16.0 <sup>c</sup> | TZ     | TZ   | TZ   | ZZ   | ZZ   |
| 1747              | 131.67                | 151.8 $\pm$ 13.1 <sup>c</sup> | ZZ     | ZZ   | ZZ   | ZZ   | TZ   |
| CK-Z <sup>a</sup> | 155.20                | 150.8 $\pm$ 14.5              | ZZ     | ZZ   | ZZ   | ZZ   | ZZ   |
| CK-T <sup>b</sup> | 196.55                | 171.3 $\pm$ 22.8 <sup>d</sup> | TT     | TT   | TT   | TT   | TT   |

ZZ, TZ, and TT indicate the Zhenshan 97 homozygotes, heterozygotes, and Teqing homozygotes at the locus of marker, respectively

<sup>a</sup> The control line with Zhenshan 97 allele of *SPP1*

<sup>b</sup> The control line with Teqing allele of *SPP1*

<sup>c</sup> No significant difference between the recombinant and CK-Z at  $P = 0.05$

<sup>d</sup> Highly significant difference between the two controls CK-Z and CK-T, and between CK-T and recombinants at  $P = 0.001$



**Table 4** Comparison of putative genes in the 107-kb region from the rice genome automated annotation database with prediction by FGENESH

| TIGR gene ID   | Putative function                                      | Homologous cDNA/EST | FGENESH gene | Comparison between TIGR gene and FGENESH gene                          | ID% <sup>a</sup> |
|----------------|--|---------------------|--------------|--|------------------|
| LOC_Os01g11990 | Expressed protein                                      | AK108579            | ORF1         | Additional sequence prior to LOC_Os01g11990                            | 81               |
| LOC_Os01g12000 | Expressed protein                                      | AK073193            | ORF2         | Missing 54 bp at the proximal end of LOC_Os01g12000                    | 52               |
| LOC_Os01g12010 | Hypothetical protein                                   | No                  | ORF3         | Same   | 100              |
| LOC_Os01g12020 | Nonspecific lipid-transfer protein precursor, putative | AK106863, AK109524  | ORF4         | Missing 12 bp at the proximal end of LOC_Os01g12020                    | 99               |
| LOC_Os01g12030 | Endoglucanase 1 precursor, putative, expressed         | AK109524            | ORF5         | Same   | 100              |
| LOC_Os01g12040 | Transposon protein, putative, mutator sub-class        | AK067920            | ORF6         | Same   | 100              |
| LOC_Os01g12060 | Hypothetical protein                                   | AK109722            | ORF7         | Same   | 100              |
| LOC_Os01g12070 | Endoglucanase 1 precursor, putative, expressed         | AK103304            | ORF8         | Same   | 100              |
| LOC_Os01g12080 | Plant-specific domain TIGR01589 family protein         | Have EST            | ORF9         | Additional 93 bp prior to LOC_Os01g12080                               | 91               |
| LOC_Os01g12090 | Expressed protein                                      | Have EST            | ORF10        | ORF10 have same start sequence but not stop sequence as LOC_Os01g12090 | 67               |
| LOC_Os01g12100 | Expressed protein                                      | AK105221            | ORF11        | Sharing the same sequence, but with slightly different coding sequence | 36               |
| LOC_Os01g12110 | Expressed protein                                      | No                  | ORF12        | Sharing the same sequence, but with slightly different coding sequence | 97               |
| LOC_Os01g12120 | Hypothetical protein                                   | No                  | ORF13        | ORF13 has same start sequence but not stop sequence as LOC_Os01g12120  | 87               |
| LOC_Os01g12130 | MtN3 protein-like, putative, expressed                 | No                  | ORF14        | Sharing the same sequence, but with slightly different coding sequence | 70               |
| LOC_Os01g12140 | Conserved hypothetical protein                         | No                  | ORF15        | Same   | 100              |
| LOC_Os01g12150 | Hypothetical protein                                   | No                  | ORF16        | Same   | 100              |
| LOC_Os01g12160 | Indole-3-acetic acid-amido synthetase GH3.3, putative  | AK072125, AK120515  | ORF17        | Sharing the same sequence, but with slightly different coding sequence | 86               |

<sup>a</sup> ID amino acid sequence identity between TIGR and FGENESH predictions

## Discussion

### A novel QTL controlling SPP

In light of the low resolution of QTL mapping in a primary population, comparison of QTL locations detected in different populations resulted in an equivocal conclusion as to whether one QTL or a QTL cluster controlling the same trait was located in the same chromosome region. Moreover, even though two linked genes were known to segregate in one primary population, it was still not easy to precisely map them because of impossible genotype inference at an individual QTL according to the target trait performance. However, with the help of advanced populations such as QTL-NILs, linked QTL could be identified separately because they can independently segregate in one NIL population. Currently, several QTL controlling the same trait have been confirmed to be tightly linked in the rice

genome, and some of them have been isolated. For example, *Hd3*, a rice heading date QTL was identified in the  $F_2$  population. Actually, two major QTL, *hd3a* and *hd3b*, which promote flowering in short-day conditions and inhibit heading in long-day conditions, respectively, were identified in the *Hd3* region in a NILs population (Monna et al. 2002a). *Xa4* and *Xa26*, two rice bacterial blight resistance genes, are tightly linked to each other in the long arm of chromosome 11 (Sun et al. 2004; Yang et al. 2003). *QTLph1* and *sd1*, which commonly control plant height, are tightly linked on chromosome 1 with a 1.7-Mb physical distance (Monna et al. 2002b; Ishimaru et al. 2004). In this study, *SPP1* was mapped to the interval of about 107 kb between RM8111 and RM490, which is closely linked to *Gn1a* and *Gn1b* controlling SPP (Ashikari et al. 2005). To genetically confirm that the SPP variation was not caused by *Gn1a* and *Gn1b*, we needed the NILs population that segregates in the region of *SPP1* but fixed in the region

covering *Gn1a* and *Gn1b*.  $F_2$ -1456 was the expected genotype in which only the *SPP1* region segregated. Its  $F_3$  family still showed a very wide variation in SPP similar to that of populations from RIL30. Hence, it is suggested that SPP variation in the population is not caused by *Gn1a* and *Gn1b* but by *SPP1*. The left flanking marker of  $F_2$ -1456, YN27, has a 1.2-Mb physical distance to *Gn1a* on the left side. The distance between *SPP1* and *Gn1a* is more than 1.2 Mb. Moreover, *Gn1b*, located in the left side of *Gn1a*, is farther from YN27 than *Gn1a* (Fig. 2). Obviously, *SPP1* is distinct from *Gn1a* and *Gn1b*.

#### Possible candidate gene

In this study, *SPP1* was fine mapped to a BAC clone. However, there are still five recombinants to fine map with new markers in the 107-kb region. The SPP could be narrowed down to a mini-region, and then its candidate gene would be determined. By contrast, it is possible to infer the candidate gene with the help of bioinformatic analysis at this time. Among the five putative known function genes, LOC\_Os01g1213 was predicted to encode MtN3 protein-like which is a transmembrane protein. The function of MtN3 family proteins is still unknown; they possibly take part in the development of root nodules and gene expressive activation of RAGs (recombination activation genes) (Gamas et al. 1996). However, *xa13*, a bacterial blight resistance gene, encodes an MtN3 family protein, which suggests a possible relationship to disease resistance (Chu et al. 2006). LOC\_Os01g12020 is a putative nonspecific LTP encoding gene. LTP can transfer lipid between different membranes and possibly participate in engendering membrane (Dansen et al. 1999). Recent studies have shown that LTP contributes to stress resistance such as cold, salt, drought, and disease resistance, and its expression can be induced by biologic and abiotic stress (Cammue et al. 1995; Gaudet et al. 2003; Wu et al. 2004). Both LOC\_Os01g12030 and LOC\_Os01g12070 are putative endoglucanase 1 precursor genes, which are involved in cell wall disassembly during organ-aging stages including fruit ripening and organ abscission (Brummell et al. 1999; Lashbrook et al. 1994). Also, endoglucanase genes are associated with disease resistance (Flors et al. 2007). Therefore, these genes seem to have nothing to do with plant inflorescence development and cannot be a candidate for *SPP1*.

It is well known that IAA is the most universal auxin in plants. It can promote cell division and growth. In rice, the endogenous level of IAA in inflorescence tissue was 10–100 times higher than that in shoot tissue, which suggests that IAA may play an important role in regulating reproductive growth and the development of inflorescent meristem (Kobayashi et al. 1989). Guan et al. (2006) used the in

vitro regeneration system of *Arabidopsis* to regenerate inflorescences when cytokinin and IAA were present. Hence, IAA is important for plant inflorescent development, which implies that IAA possibly regulates SSP in rice. In *Arabidopsis thaliana*, the *pin-formed* (*pin*) mutant had no flowers or had abnormal flowers on the inflorescence axis (Goto et al. 1991; Okada et al. 1991; Okada and Shimura 1994). The concentration of IAA and the activity of auxin polar transport in the inflorescence axis of the *pin* mutant were much lower than those of the wild-type (Enkheim ecotype) (Okada et al. 1991). Moreover, the application of inhibitors of IAA polar transport to the wild type plant of *Arabidopsis* induced the formation of an inflorescence axis similar to that of the *pin-formed* mutant (Okada et al. 1991; Okada and Shimura 1994). In addition, other evidence also showed that normal concentration of endogenous IAA and/or the activity of IAA polar transport is essential for the differentiation and formation of the floral meristem (Oka et al. 1999; Aloni et al. 2006). Hence, the LOC\_Os01g12160, which codes IAA synthetase, is considered the most interesting candidate gene. Paying more attention to confirming the candidate gene function by transformation or by checking its mutation will be worthwhile.

#### Potential exploitation in rice yield improvement

Recently, several research groups have cloned the genes related to grain yield in rice. According to their results, the genes can be classified into two types: high grain yield accompanied by elongation of life cycle (type I) and high grain yield without a change in life cycle (type II). For example, *Ghd7* could produce more rice yield by delaying flowering time accompanied by an increase in plant height (Xue et al. 2008). However, yield in the *Gn1a* plant increased 40 grains per main panicle without a change in flowering time and plant height. Heading date is an important factor to determine seasonal and regional adaptation of rice. In terms of gene characterization, the rice growing region should be considered when exploiting type I genes for rice production. Otherwise, the plant cannot take full advantage of light and temperature for rice production, or rice cannot grow normally during the flowering period because of low temperature. Both cases would cause a large loss in yield. In a word, the exploitation of type I genes is dependent on rice growing regions and seasons. For type II genes, their exploitation has a large degree of freedom. Like *SPP1* reported in this study, it had no effect on seed setting and 1,000-grain weight (no QTL was detected in the NILs- $F_2$  population), but it had a major effect on SPP. Positive *SPP1* homozygotes could result in a about 10-g yield increase per plant compared with negative homozygotes. The positive allele of *SPP1*, theoretically, can be transferred to

rice varieties grown in any rice regions. Hence, type II gene is more favorable for rice breeding. In addition, *SPP1* showed dominant effects on SPP and grain yield, which indicated that the Teqing homozygote has more yield than the heterozygote. Hence, in theory, *SPP1* should be simultaneously transferred to both female and male parents to produce a better hybrid.

**Acknowledgments** This work was supported by a grant from the National Key Program on Basic Research and Development of China, a grant from National Natural Science Foundation of China, and a grant for New Century Excellent Talents from the Education Ministry of China.

## References

- Aloni R, Aloni E, Langhans M, Ullrich CI (2006) Role of auxin in regulating *Arabidopsis* flower development. *Planta* 223:315–328
- Ashikari M, Sakakibara H, Lin S, Yamamoto T, Takashi T, Nishimura A, Angeles ER, Qian Q, Kitano H, Matsuoka M (2005) Cytokinin oxidase regulates rice grain production. *Science* 309:741–745
- Brummell DA, Hall B, Bennett AB (1999) Antisense suppression of tomato endo-1, 4-beta glucanase Cel2 mRNA accumulation increases the force required to break fruit abscission zones but does not affect fruit softening. *Plant Mol Biol* 40:615–622
- Cammue B, Thevissen K, Hendriks M, Kristel Eggermont K, Goderis I, Proost P, Damme J, Osborn R, Guerbette F, Kader J, Broekaert W (1995) A potent antimicrobial protein from onion seeds showing sequence homology to plant lipid transfer proteins. *Plant Physiol* 109:445–455
- Chu Z, Yuan M, Yao J, Ge X, Yuan B, Xu C, Li X, Fu B, Li Z, Benetzen JL, Zhang Q, Wang S (2006) Promoter mutations of an essential gene for pollen development result in disease resistance in rice. *Genes Dev* 20:1250–1255
- Dansen T, Westerman J, Wouters F, Wanders R, Hoek A, Gadella T, Wirtz K (1999) High-affinity binding of very-long-chain fatty acyl-CoA esters to the peroxisomal non-specific lipid-transfer protein (sterol carrier protein22). *Biochem J* 339:193–199
- Fan CC, Xing YZ, Mao HL, Lu TT, Han B, Xu CG, Li XH, Zhang QF (2006) *GS3*, a major QTL for grain length and weight and minor QTL for grain width and thickness in rice, encodes a putative transmembrane protein. *Theor Appl Genet* 112:1164–1171
- Flors V, Leyva M, Vicedo B, Finiti I, Real M, Garcia-Agustín P, Bennett B, González-Bosch C (2007) Absence of the endo-b-1, 4-glucanases Cel1 and Cel2 reduces susceptibility to *Botrytis cinerea* in tomato. *Plant J* 52:1027–1040
- Gamas P, Niebel FC, Lescure N, Cullimore JV (1996) Use of a subtractive approach to identify new *Medicago truncatula* genes induced during root nodule development. *Mol Plant Microbe Interact* 9:233–242
- Gaudet D, Laroche A, Frick M, Huel R, Puchalski B (2003) Cold induced expression of plant defensin and lipid transfer protein transcripts in winter wheat. *Physiol Plantarum* 117:1399–3054
- Goto N, Katoh N, Kranz AR (1991) Morphogenesis of floral organs in *Arabidopsis*: predominant carpel formation of the *pin-formed* mutant. *Jpn J Genet* 66:551–567
- Guan CM, Zhu SS, Li XG, Zhang XS (2006) Hormone-regulated inflorescence induction and *TFL1* expression in *Arabidopsis* callus in vitro. *Plant Cell Rep* 25:1133–1137
- Hittalmani S, Huang N, Courtois B, Venuprasad R, Shashidhar HE, Zhuang JY, Zheng KL, Liu GF, Wang GC, Sidhu JS, Srivantaneeyakul S, Singh VP, Bagali PG, Prasanna HC, McLaren G, Khush GS (2003) Identification of QTL for growth- and grain yield-related traits in rice across nine locations of Asia. *Theor Appl Genet* 107:679–690
- Ishimaru K, Ono K, Kashiwagi T (2004) Identification of a new gene controlling plant height in rice using the candidate-gene strategy. *Planta* 218:388–395
- Kobayashi M, Sakurai A, Saka H, Takahashi N (1989) Fluctuation of the endogenous IAA level in rice during its life cycle. *Agric Biol Chem* 53:1089–1094
- Lander ES, Botstein D (1989) Mapping Mendelian factors underlying quantitative traits using RFLP linkage maps. *Genetics* 121:185–199
- Lashbrook CC, González-Bosch C, Bennett AB (1994) Two divergent endo-beta-1, 4-glucanase genes exhibit overlapping expression in ripening fruit and abscising flowers. *Plant Cell* 6:1485–1493
- Li Z, Pinson SRM, Park WD, Paterson AH, Stansel JW (1997) Epistasis for three grain yield components in rice (*Oryza sativa* L.). *Genetics* 145:453–465
- Li JM, Thomson M, McCouch RS (2004) Fine mapping of a grain weight quantitative trait locus in the pericentromeric region of rice chromosome 3. *Genetics* 168:2187–2195
- Lincoln S, Daley M, Lander E (1992) Constructing genetic maps with MAPMAKER/EXP 3.0. In: Whitehead Institute technical report, 3rd edn. Whitehead Institute, Cambridge
- Lincoln SE, Daly MJ, Lander ES (1993) Mapping genes controlling quantitative traits with MAPMAKER/QTL1.1: a tutorial and reference manual, 2nd edn. Whitehead Institute, Cambridge
- McCouch SR, Teytelman L, Xu Y, Lobos KB, Clare K, Walton M, Fu B, Maghirang R, Li Z, Xing Y, Zhang Q, Kono I, Yano M, Fjellstrom R, DeClerck G, Schneider D, Cartinhour S, Ware D, Stein L (2002) Development and mapping of 2, 240 new SSR markers for rice (*Oryza sativa* L.). *DNA Res* 9:199–207
- Monna L, Lin X, Kojima S, Sasaki T, Yano M (2002a) Genetic dissection of a genomic region for a quantitative trait locus, *Hd3*, into two loci, *Hd3a* and *Hd3b*, controlling heading date in rice. *Theor Appl Genet* 104:772–778
- Monna L, Kitazawa N, Yoshino R, Suzuki J, Masuda H, Maehara Y, Tanji M, Sato M, Nasu S, Minobe Y (2002b) Positional cloning of rice semidwarfing gene, *sd-1*: rice “green revolution gene” encodes a mutant enzyme involved in gibberellin synthesis. *DNA Res* 9:11–17
- Murray MG, Thompson WF (1980) Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Res* 8:4321
- Oka M, Miyamoto K, Okada K, Ueda J (1999) Auxin polar transport and flower formation in *Arabidopsis thaliana* transformed with indoleacetamide hydrolase (*iaaH*) gene. *Plant Cell Physiol* 40:231–237
- Okada K, Shimura Y (1994) Genetic analyses of signalling in flower development using *Arabidopsis*. *Plant Mol Biol* 26:1357–1377
- Okada K, Ueda J, Komaki MK, Bell CJ, Shimura Y (1991) Requirement of the auxin polar transport system in early stages of *Arabidopsis* floral bud formation. *Plant Cell* 3:677–684
- Ren ZH, Gao JP, Li LG, Cai XL, Huang W, Chao DY, Zhu MZ, Wang ZY, Luan S, Lin HX (2005) A rice quantitative trait locus for salt tolerance encodes a sodium transporter. *Nat Genet* 37:1141–1146
- Septiningsih EM, Prasetyono J, Lubis E, Tai TH, Tjubaryat T, Moeljopawiro S, McCouch SR (2003) Identification of quantitative trait loci for yield and yield components in an advanced backcross population derived from the *Oryza sativa* variety IR64 and the wild relative *O. rufipogon*. *Theor Appl Genet* 107:1419–1432
- Song XJ, Huang W, Shi M, Zhu MZ, Lin HX (2007) A QTL for rice grain width and weight encodes a previously unknown RING-type E3 ubiquitin ligase. *Nat Genet* 39:623–630
- Sun X, Cao Y, Yang Z, Xu C, Li X, Wang S, Zhang Q (2004) *Xa26*, a gene conferring resistance to *Xanthomonas oryzae* pv. *oryzae* in rice, encodes an LRR receptor kinase-like protein. *Plant J* 37:517–527



- Temnykh S, Park WD, Ayres N, Cartinhour S, Hauck N, Lipovich L, Cho YG, Ishii T, McCouch SR (2000) Mapping and genome organization of microsatellite sequences in rice (*Oryza sativa* L.). *Theor Appl Genet* 100:697–712
- Temnykh S, Declerck G, Luashova A, Lipovich L, Cartinhour S, McCouch S (2001) Computational and experimental analysis of microsatellites in rice (*Oryza sativa* L.): frequency, length variation, transposon associations, and genetic marker potential. *Genome Res* 11:1441–1452
- Tian F, Zhu Z, Zhang B, Tan L, Fu Y, Wang X, Sun CQ (2006) Fine mapping of a quantitative trait locus for grain number per panicle from wild rice (*Oryza ruffipogon* Griff.). *Theor Appl Genet* 113:619–629
- Tuinstra MR, Ejeta G, Goldsbrough PB (1997) Heterogeneous inbred family (HIF) analysis: a method for developing near-isogenic lines that differ at quantitative trait loci. *Theor Appl Genet* 95:1005–1011
- Wu KS, Tanksley SD (1993) Abundance, polymorphism and genetic mapping of microsatellites in rice. *Mol Gen Genet* 241:225–235
- Wu G, Robertson AJ, Liu X, Zheng P, Wilen RW, Nesbitt NT, Gusta LV (2004) A lipid transfer protein gene *BG-14* is differentially regulated by abiotic stress, ABA, anisomycin, and sphingosine in bromegrass (*Bromus inermis*). *Plant Physiol* 161:449–458
- Xing YZ, Tan YF, Hua JP, Sun XL, Xu CG, Zhang Q (2002) Characterization of the main effects, epistatic effects and their environmental interactions of QTLs on the genetic basis of yield traits in rice. *Theor Appl Genet* 105:248–257
- Xing YZ, Tang WJ, Xue WY, Xu CG, Zhang Q (2008) Fine mapping of a major quantitative trait loci, *qSSP7*, controlling the number of spikelets per panicle as a single Mendelian factor in rice. *Theor Appl Genet* 116:789–796
- Xue W, Xing Y, Weng X, Zhao Y, Tang W, Wang L, Zhou H, Yu S, Xu C, Li X, Zhang Q (2008) Natural variation in *Ghd7* is an important regulator of heading date and yield potential in rice. *Nat Genet* 40(6):761–767
- Yamamoto T, Kuboki Y, Lin SY, Sasaki T, Yano M (1998) Fine mapping of quantitative trait loci *Hd-1*, *Hd-2* and *Hd-3*, controlling heading date of rice, as single Mendelian factors. *Theor Appl Genet* 97:37–44
- Yamamoto T, Lin HX, Sasaki T, Yano M (2000) Identification of heading date quantitative trait locus *Hd6* and characterization of its epistatic interactions with *Hd2* in rice using advanced back-cross progeny. *Genetics* 154:885–891
- Yang Z, Sun X, Wang S, Zhang Q (2003) Genetic and physical mapping of a new gene for bacterial blight resistance in rice. *Theor Appl Genet* 106(8):1467–1472
- Yu SB, Li JX, Xu CG, Tan YF, Gao YJ, Li XH, Zhang QF (1997) Importance of epistasis as the genetic basis of heterosis in an elite rice hybrid. *Proc Natl Acad Sci USA* 94:9226–9231
- Zhang YS, Luo LJ, Xu CG, Zhang QF, Xing YZ (2006) Quantitative trait loci for panicle size, heading date and plant height co-segregating in trait-performance derived near-isogenic lines of rice (*Oryza sativa*). *Theor Appl Genet* 113:361–368
- Zhuang JY, Lin HX, Lu J, Qian HR, Hittalmani S, Huang N, Zheng KL (1997) Analysis of QTL  $\times$  environment interaction for yield components and plant height in rice. *Theor Appl Genet* 95:799–808
- Zhuang JY, Fan YY, Rao ZM, Wu JL, Xia YW, Zheng KL (2002) Analysis on additive effects and additive-by-additive epistatic effects of QTLs for yield traits in a recombinant inbred line population of rice. *Theor Appl Genet* 105:1137–1145