

## DIRICHLET PRINCIPLE OF HARMONIC MAPS

ZUJIN ZHANG

ABSTRACT. We define harmonic maps as the critical point of Dirichlet energy functional. This is [1, 1.1].

Let

1.  $(M^n, g)$  be a Riemannian manifold with or without boundary;
2.  $(N^l, h)$  be a compact Riemannian manifold without boundary (closed).

The we may define the **Dirichlet energy functional**

$$E(u) = \int_M e(u) dv_g,$$

where  $e(u)$  is the **Dirichlet energy density function**, the expression of which in local coordinates  $(U, x^\alpha), (V, u^i)$  is

$$e(u) \equiv \frac{1}{2} |\nabla u|_g^2 = \frac{1}{2} g^{\alpha\beta}(x) h_{ij}(u(x)) \frac{\partial u^i}{\partial x^\alpha} \frac{\partial u^j}{\partial x^\beta}.$$

**Definition 1.** A map  $u \in C^2(M, N)$  is a **harmonic map** if it is a critical point of the Dirichlet energy functional  $E$ .

**Proposition 2.** A map  $u \in C^2(M; N)$  is a **harmonic map** iff  $u$  satisfies

$$\Delta_g u^i + g^{\alpha\beta} \Gamma_{jk}^i(u) \frac{\partial u^j}{\partial x^\alpha} \frac{\partial u^k}{\partial x^\beta} = 0, \text{ in } M, 1 \leq i \leq l.$$

Here

---

*Key words and phrases.* harmonic map, Dirichlet principle.

1.  $\Delta_g$  is the **Laplace-Beltrami operator** on  $(M, g)$  given by

$$\Delta_g = \frac{1}{\sqrt{g}} \frac{\partial}{\partial x^\alpha} \left( \sqrt{g} g^{\alpha\beta} \frac{\partial}{\partial x^\beta} \right);$$

2. and  $\Gamma_{jk}^i$  is the **Christoffel symbol** of the metric  $h$  on  $N$  given by

$$\Gamma_{jk}^i = \frac{1}{2} h^{il} (h_{li,k} + h_{kl,j} - h_{jk,l}).$$

*Proof.* 1. Let  $U \subset M$  be any coordinate chart and  $\varphi \in C_0^2(U; \mathbf{R}^l)$ .

Then we have

$$\begin{aligned} 0 &= \frac{d}{dt} \Big|_{t=0} \left[ \frac{1}{2} \int_M g^{\alpha\beta} h_{ij}(u + t\varphi) (u_\alpha^i + t\varphi_\alpha^i) (u_\beta^j + t\varphi_\beta^j) \sqrt{g} dx \right] \\ &= \frac{1}{2} \int_M g^{\alpha\beta} h_{ij,k}(u) \varphi_k u_\alpha^i u_\beta^j \sqrt{g} dx + \int_M g^{\alpha\beta} h_{ij}(u) u_\alpha^i \varphi_\beta^j \sqrt{g} dx. \end{aligned}$$

2. Direct computations show

$$\begin{aligned} \int_M \Delta_g u^i h_{ij}(u) \varphi^j dv_g &= \int_M \frac{\partial}{\partial x^\alpha} \left( \sqrt{g} g^{\alpha\beta} \frac{\partial u^i}{\partial x^\beta} \right) h_{ij}(u) \varphi^j dx \\ &= - \int_M \sqrt{g} g^{\alpha\beta} u_\beta^i (h_{ij,k}(u) u_\alpha^k \varphi^j + h_{ij}(u) \varphi_\alpha^j) dx \\ &= - \frac{1}{2} \int_M \sqrt{g} g^{\alpha\beta} u_\alpha^i u_\beta^j (h_{ik,j} + h_{kj,i} - h_{ij,k}) (u) \varphi^k dx \\ &= - \int_M g^{\alpha\beta} \Gamma_{ij}^l(u) h_{lk}(u) u_\alpha^i u_\beta^j \varphi^k dv_g. \end{aligned}$$

This implies that

$$\Delta u^i + g^{\alpha\beta} \Gamma_{kl}^i(u) u_\alpha^k u_\beta^l = 0, \quad \forall 1 \leq i \leq l.$$

□

## REFERENCES

- [1] F.H. Lin, C.Y. Wang, *The analysis of harmonic maps and their heat flows*. World Scientific Publishing Co. Pte. Ltd. 2008.

DEPARTMENT OF MATHEMATICS, SUN YAT-SEN UNIVERSITY, GUANGZHOU,  
510275, P.R. CHINA

*E-mail address:* uia.china@gmail.com